

# AQA Further Pure 1

## Complex Numbers

### Section 1: Introduction to Complex Numbers

#### Notes and Examples

These notes contain subsections on

- The number system
- Adding and subtracting complex numbers
- Multiplying complex numbers
- Complex conjugates
- Equations with complex roots
- Equating real and imaginary parts

#### The number system

In your learning of mathematics, you have come across different types of number at different stages. Each time you were introduced to a new set of numbers, this allowed you to solve a wider range of problems.


The first numbers that you came across were the counting numbers (natural numbers). These allowed you to solve equations like  $x + 2 = 5$ .

Later you would meet negative numbers, which allowed you to solve equations like  $x + 5 = 2$ , and rational numbers, which meant you could solve equations like  $2x = 5$ .

When irrational numbers were included, you could solve equations like  $x^2 = 2$ .

However, there are still equations which you cannot solve, such as  $x^2 = -4$ . You know that there are no real numbers which satisfy this equation. However, this equation, and others like it, can be solved using imaginary numbers, which are based on the number  $i$ , which is defined as  $\sqrt{-1}$ .

The diagram below shows the relationships between different types of number.



# AQA Further Pure 1

This type of diagram is called a Venn diagram (you may have met these before if you have studied any Statistics) and it shows the relationships between sets, in this case sets of numbers. This diagram deals with the real numbers, which include all numbers which you have come across until now. Notice that the positive and negative integers (whole numbers) are subsets of the rational numbers. This means that all integers are also rational numbers, but there are other rational numbers which are not integers, such as  $\frac{3}{2}$  or  $-\frac{7}{11}$ . Similarly, all rational numbers are real numbers, but there are other real numbers which are not rational, such as  $\sqrt{3}$  and  $\pi$ .

In this chapter you will see that the real numbers are also a subset of a larger set called the complex numbers. You will be looking at numbers which lie outside the set of real numbers. Complex numbers involve both real and imaginary numbers.

## Adding and subtracting complex numbers

To add two complex numbers, you need to add the real parts and add the imaginary parts. Similarly, to subtract one complex number from another, deal with the real and imaginary parts separately.



### Example 1

The complex numbers  $z$  and  $w$  are given by

$$z = 3 + 2i$$

$$w = 1 - 4i$$

Find:

(i)  $z + w$

(ii)  $z - w$

(iii)  $w - z$



### Solution

$$\begin{aligned} \text{(i) } z + w &= (3 + 2i) + (1 - 4i) \\ &= (3 + 1) + (2i - 4i) \\ &= 4 - 2i \end{aligned}$$

Add the real parts and  
add the imaginary parts

$$\begin{aligned} \text{(ii) } z - w &= (3 + 2i) - (1 - 4i) \\ &= (3 - 1) + (2i + 4i) \\ &= 2 + 6i \end{aligned}$$

Subtract the real parts and  
subtract the imaginary parts

$$\begin{aligned} \text{(iii) } w - z &= (1 - 4i) - (3 + 2i) \\ &= (1 - 3) + (-4i - 2i) \\ &= -2 - 6i \end{aligned}$$



For practice in examples like the one above, try the interactive resource [Addition and subtraction of complex numbers.](#)

# AQA Further Pure 1

## Multiplying complex numbers

Multiplication of two complex numbers is similar to multiplying out a pair of brackets. Each term in the first bracket must be multiplied by each term in the second bracket. You can then simplify, remembering that  $i^2 = -1$ .



### Example 2

Find

- (i)  $(3 + 4i)(2 + i)$
- (ii)  $(2 - i)(3 + 2i)$
- (iii)  $(2 + 3i)(2 - 3i)$

### Solution

- (i)  $(3 + 4i)(2 + i) = 6 + 3i + 8i + 4i^2$   
 $= 6 + 11i - 4$   
 $= 2 + 11i$
- (ii)  $(4 - i)(3 + 2i) = 12 + 8i - 3i - 2i^2$   
 $= 12 + 5i + 2$   
 $= 14 + 5i$
- (iii)  $(2 + 3i)(2 - 3i) = 4 - 6i + 6i - 9i^2$   
 $= 4 + 9$   
 $= 13$

Multiply out the brackets

Using  $i^2 = -1$



For practice in examples like the one above, try the interactive resource [Multiplication of complex numbers](#).

## Complex conjugates

In part (iii) of Example 2, the result of multiplying two complex numbers is a real number. This is always the case when the complex number  $a + bi$  is multiplied by the complex number  $a - bi$ . The complex number  $a - bi$  is called the **complex conjugate** of  $a + bi$ . For any complex number  $z$ , the complex conjugate is written as  $\bar{z}$  or  $z^*$ .



For practice in complex conjugates, try the interactive resource [Conjugate of a complex number](#).

## Equations with complex roots

When you first learned to solve quadratic equations using the quadratic formula, you found that some quadratic equation had no real solutions. However, using complex numbers you can find solve all quadratic equations.

# AQA Further Pure 1



## Example 3

Solve the quadratic equation

$$x^2 + 6x + 13 = 0$$

## Solution

Using the quadratic formula with  $a = 1$ ,  $b = 6$ ,  $c = 13$

$$\begin{aligned}x &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\&= \frac{-6 \pm \sqrt{36 - 4 \times 1 \times 13}}{2 \times 1} \\&= \frac{-6 \pm \sqrt{-16}}{2} \\&= \frac{-6 \pm 4i}{2} \\&= -3 \pm 2i\end{aligned}$$

The solutions of the equation are  $x = -3 + 2i$  and  $x = -3 - 2i$

Notice that the quadratic equation in Example 3 has two complex solutions which are a pair of complex conjugates. All quadratic equations with real coefficients have two solutions: either two real solutions (which could be a repeated solution) or two complex solutions which are a pair of complex conjugates.

The next example shows how you can find a quadratic equation with roots at particular complex values. A quadratic equation with roots at  $x = a$  and  $x = b$  can be written as  $(x - a)(x - b) = 0$ , and this also applies to situations where the roots are complex numbers. Alternatively, you can consider the sum and the products of the roots. Both approaches are shown in Example 4.



## Example 4

Find the quadratic equation which has roots at  $x = 4 + 2i$  and  $x = 4 - 2i$ .

## Solution 1

$$\begin{aligned}(x - (4 + 2i))(x - (4 - 2i)) &= 0 \\(x - 4 - 2i)(x - 4 + 2i) &= 0 \\(x - 4)^2 - (2i)^2 &= 0 \\x^2 - 8x + 16 + 4 &= 0 \\x^2 - 8x + 20 &= 0\end{aligned}$$

The two middle terms cancel out since this expression is of the form  $(x - a)(x + a)$

## Solution 2

The sum of the roots is  $4 + 2i + 4 - 2i = 8 \Rightarrow -\frac{b}{a} = 8$

The product of the roots is  $(4 + 2i)(4 - 2i) = 16 + 4 = 20 \Rightarrow \frac{c}{a} = 20$

Taking  $a = 1$  gives  $b = -8$  and  $c = 20$

The quadratic equation is  $x^2 - 8x + 20 = 0$ .

# AQA Further Pure 1



The Flash resource [Working with complex numbers](#) tests you on multiplication, complex conjugates and equations with complex roots.



You can also try the Complex numbers Hexagonal puzzle.



Using complex numbers, you have seen that all quadratic equations have two solutions. In a similar way, all cubic equations have three solutions, all quartic equations have four solutions, and so on. For all polynomial equations with real coefficients, any complex roots always occur in conjugate pairs. So the three roots of a cubic equation could be three real roots (possibly including repeated roots) or one real root and a conjugate pair of complex roots. Similarly, the four roots of a quartic equation could be four real roots (possibly including repeated roots), two real roots and a conjugate pair of complex roots, or two conjugate pairs of complex roots.

However, in Further Pure 1 you will only be required to solve quadratic equations with complex roots.

## Equating real and imaginary parts

For two complex numbers to be equal, then the real parts must be equal and the imaginary parts must be equal. So one equation involving complex numbers can be written as two equations, one for the real parts, one for the imaginary parts.

The examples below show how this technique can be used to solve equations involving complex numbers.



### Example 5

Solve the equation  $3z - 2z^* = 4 - 15i$

### Solution

Let  $z = x + iy$

$$3z - 2z^* = 4 - 15i$$

$$3(x + iy) - 2(x - iy) = 4 - 15i$$

$$3x + 3iy - 2x + 2iy = 4 - 15i$$

$$x + 5iy = 4 - 15i$$

Equating real parts:  $x = 4$

Equating imaginary parts:  $5y = -15 \Rightarrow y = -3$

$$z = 4 - 3i$$



# AQA Further Pure 1

In Example 5 the values of  $x$  and  $y$  could be found immediately by equating real and imaginary parts. In the next example, equating real and imaginary parts gives you a pair of linear simultaneous equations to solve.



## Example 6

Solve the equation

$$(3 - 2i)(z - 1 + 4i) = 7 + 4i$$

## Solution

Let  $z = x + iy$

$$(3 - 2i)(x + iy - 1 + 4i) = 7 + 4i$$

$$(3 - 2i)((x - 1) + i(y + 4)) = 7 + 4i$$

$$3(x - 1) - 2i(x - 1) + 3i(y + 4) - 2i^2(y + 4) = 7 + 4i$$

$$3(x - 1) - 2i(x - 1) + 3i(y + 4) + 2(y + 4) = 7 + 4i$$

$$\text{Equating real parts:} \quad 3(x - 1) + 2(y + 4) = 7 \quad \Rightarrow 3x + 2y = 2 \quad \textcircled{1}$$

$$\text{Equating imaginary parts:} \quad -2(x - 1) + 3(y + 4) = 4 \quad \Rightarrow -2x + 3y = -10 \quad \textcircled{2}$$

$$\textcircled{1} \times 2 \quad 6x + 4y = 4$$

$$\textcircled{2} \times 3 \quad -6x + 9y = -30$$

$$\text{Adding:} \quad 13y = -26$$

$$y = -2, x = 2$$

$$z = 2 - 2i$$



## Example 7

Find real numbers  $a$  and  $b$  with  $a > 0$  such that  $(a + bi)^2 = 16 - 30i$ .

## Solution

$$(a + bi)^2 = 16 - 30i$$

$$a^2 + 2abi + b^2i^2 = 16 - 30i$$

$$a^2 + 2abi - b^2 = 16 - 30i$$

$$\text{Equating imaginary parts:} \quad 2ab = -30 \Rightarrow b = -\frac{15}{a}$$

$$\text{Equating real parts:} \quad a^2 - b^2 = 16$$

$$\text{Substituting:} \quad a^2 - \frac{225}{a^2} = 16$$

$$\text{Multiplying through by } a^2: \quad a^4 - 225 = 16a^2$$

$$a^4 - 16a^2 - 225 = 0$$

This is a quadratic in  $a^2$  and can be factorised:

$$(a^2 - 25)(a^2 + 9) = 0$$

Since  $a$  is real,  $a^2 + 9$  cannot be equal to zero.

Since  $a > 0$ , the only possible solution is  $a = 5$ .

$$b = -\frac{15}{a} \Rightarrow b = -\frac{15}{5} = -3$$

$$a = 5, b = -3$$



## AQA Further Pure 1

Notice that in this example you are finding the square root of a complex number. The square root of  $16 - 30i$  is  $5 - 3i$ . You can find the other square root of this complex number by allowing  $a$  to be negative, so  $a = -5$  and then  $b = 3$ . So the other square root is  $-5 + 3i$ . As with real numbers, one square root is the negative of the other.